arXiv:cond-mat/0512418v1 [cond-mat.stat-mech] 17 Dec 2005

Watanabe and Hu Reply: In a recent Letter [1], Watanabe, Yukawa, Ito, and Hu (WYIH) proposed the idea of superscaling for percolation on rectangular domains and used numerical data near the effective critical density ρ_c' to support such idea. In [2], Pruessner and Moloney (PM) pointed out some problems in [1]. Most of these problems are caused by the differences in using ρ_c' or the bulk critical density ρ_c in the analysis. Figure 2 of [1] shows clearly that at $\rho = \rho_c$ the existence probabilities E_p for different aspect ratios R are different [3]. Thus it is natural that we consider a horizontal line at $E_p = 0.5$ in Fig. 2(a) to define the effective critical density ρ_c' for curves with different aspect ratios R and using data near ρ_c' to study superscaling shown in Fig. 4 of [1]. Please note that at ρ_c', the correlation length is of the order of vertical linear dimension L. By considering E_p near ρ_c' rather than near ρ_c, we will not have the problems mentioned at the second and the third paragraphs of [2]. To avoid the confusion, in the abstract and the first paragraph of [1] we should replace "deviation from the critical point ϵ" by "deviation from the effective critical point ρ_c', ϵ,"; we should also replace the term "the critical point" below Eq. (1) and "ρ_c" of Eq. (2) by "the effective critical point ρ_c'" and "ρ_c'", respectively.

The value of b in Eq. (10) of [1] was determined by PM to be $3/8(\equiv y_t/2)$ based on $P(L, R, \epsilon) = P(RL, R^{-1}, \epsilon)$, while WYIH obtained $b = 0.05$. This discrepancy is caused by differences in definitions of percolating clusters. PM used only the largest cluster to define the percolation probability P and thus obtained $P(L, R, \epsilon) = P(RL, R^{-1}, \epsilon)$. WYIH used all clusters extending from the top to the bottom of the system to define P and have $P(L, R, \epsilon) \geq P(RL, R^{-1}, \epsilon)$ for $R > 1$. Besides, WYIH used the effective critical density ρ_c' in Fig. 7 of [1] to determine $b = 0.05$, while PM did not used ρ_c'. To avoid the confusion, we should also use the effective critical density rather than the critical density to define t in Eq. (10) of [1] as in the case of Fig. 7.

Since a horizontal line at $E_p = E_p(L, 0, 1)$ in Fig. 2 of [1] has been used to defined ρ_c', Fig. 2 indicates that ρ_c' depends on L and R and $\lim_{L\to\infty} E_p(L, \rho_c' - \rho_c, R) - E_p(L, 0, R) \neq 0$ for $R \neq 1$. Thus PM correctly pointed out that the difference between using ρ_c' and ρ_c does not vanish in the limit $L \to \infty$ for $R \neq 1$. For this reason, we should replace "correct for finite-size effects" above Eq. (9) in [1] by "study superscaling behavior".

The meaning of the *nonlinear scaling variable* questioned by PM can be understood from the mapping function $f(\rho)$ defined by $E_p(L, \rho, R) = E_p(L, f(\rho), 1)$, with f determined numerically and plotted in Fig. 1 which shows nonlinear dependence on ρ for $R > 1$ and is consistent with Fig. 6 and Table I of [1]. Finite-size scaling implies $E_p(L, f(\rho), 1) \sim F((f(\rho) - \rho_c)L^{y_t})$; combining this with the expansion of $f(\rho)$ at ρ_c' up to the linear term with $(\partial f/\partial\rho)_{\rho_c'} \sim R^a$ gives $E_p(L, \rho, R) \sim F((\rho - \rho_c')L^{y_t}R^a)$, which shows superscaling of E_p with scaling variable $x = (\rho - \rho_c')L^{y_t}R^a$ and ρ_c' (instead of ρ_c) enters x naturally. We find that $\ln(\partial f/\partial\rho)_{\rho_c'}$ increases linearly with $\ln R$ with a well defined exponent $a = 0.14$ [4] which is consistent with that obtained in [1]. Similar calculations of the mapping functions of the percolation probability give the exponent $b = 0.05$ [4] which is also consistent with that in [1].



FIG. 1: The mapping functions f in Eq. (1), which are nonlinear for $R > 1$. The inset shows how to obtain f from E_p. The solid circles are the effective critical points ρ_c'.

Scaling can be considered as "data collapse" for critical systems with different sizes [5]. We found very good data collapse for systems with different aspect ratios in [1]. It is appropriate to call such results "superscaling".

This work was supported by Grants NSC 93-2112-M 001-027 and NSC 94-2119-M-002-001.

Hiroshi Watanabe[1] and Chin-Kun Hu[2]

[1]Department of Complex Systems Science, Graduate School of Information Science, Nagoya University, Furou-chou, Chikusa-ku, Nagoya 464-8601, Japan

[2]Institute of Physics, Academia Sinica, Nankang, Taipei 11529, Taiwan

Received 10 June 2005
PACS numbers 05.50.+q, 24.10.Lx, 89.75.Da

[1] H. Watanabe, S. Yukawa, N. Ito, and C.-K. Hu, Phys. Rev. Lett. **93**, 19601 (2004).
[2] G. Pruessner and N. R. Moloney, Phys. Rev. Lett. **95**, 258901 (2005).
[3] J. L. Cardy, J. Phys. A **25**, L201 (1992).
[4] H. Watanabe and C.-K. Hu, unpublished data.
[5] H. E. Stanley, Rev. Mod. Phys. **71**, S358 (1999).